Check for updates

Received: 17 July 2023 | Accepted: 15 March 2024

DOI: 10.1111/mms.13125

ARTICLE

Marine Mammal Science

Exposure of satellite tagged bowhead whales (*Balaena mysticetus*) to transiting vessels in the Eastern Canadian Arctic

Morgan J. Martin[1,2,3] | William D. Halliday[1,4] | Steven H. Ferguson[5] | Brent G. Young[5] | Rachel Charish[1] | Jackie Dawson[6] | Sarah M. E. Fortune[7] | Francis Juanes[2] | Levi Qaunaq[8] | Stephen J. Insley[1,2]

[1]Wildlife Conservation Society Canada, Whitehorse, Yukon, Canada

[2]Department of Biology, University of Victoria, Victoria, British Columbia, Canada

[3]Center for Marine Acoustics, Bureau of Ocean Energy Management, Virginia

[4]School of Earth and Ocean Sciences, University of Victoria, Victoria, British Columbia, Canada

[5]Fisheries and Oceans Canada, Winnipeg, Manitoba, Canada

[6]Department of Geography, Environment, and Geomatics, University of Ottawa, Ottawa, Ontario, Canada

[7]Department of Oceanography, Dalhousie University, Halifax, Nova Scotia, Canada

[8]Community member, Igloolik, Nunavut, Canada

Correspondence
Morgan J. Martin, Center for Marine Acoustics, Bureau of Ocean Energy Management, 45600 Woodland Road, Sterling, VA 20166.
Email: mjmartin@sandiego.edu

Abstract

Climate change poses new challenges to Arctic marine mammals, with increasing vessel traffic and associated underwater noise pollution emerging as significant threats. The bowhead whale (*Balaena mysticetus*), an endemic Arctic cetacean, faces these new threats. The Eastern Canada-West Greenland (ECWG) bowhead whale population migrates through areas with the highest levels of vessel traffic in the Canadian Arctic. Here, we document the spatial and temporal overlap between 36 satellite-tagged ECWG bowhead whales and vessels equipped with Automatic Identification System (AIS) transponders during 2012–2017. We report 1,145 instances where vessels were within 125 km of a tagged whale, with 306 occurrences within distances ≤50 km. Overlap between vessels and tagged bowhead whales was quantified monthly within years to investigate individual whale encounter rates. Results indicate that ECWG bowhead whales encounter the majority (79%) of vessels annually during August–October,

This is an open access article under the terms of the Creative Commons Attribution-NonCommercial-NoDerivs License, which permits use and distribution in any medium, provided the original work is properly cited, the use is non-commercial and no modifications or adaptations are made.
© 2024 The Authors. *Marine Mammal Science* published by Wiley Periodicals LLC on behalf of Society for Marine Mammalogy.

Funding information
Nunavut Wildlife Research Trust Fund; ArcticNet Centre of Excellence, Grant/Award Number: awarded to Steve Ferguson; Weston Family Foundation; Canada Nature Fund for Aquatic Species at Risk; Fisheries and Oceans Canada; Mitacs, Grant/Award Number: Accelerate Program; MEOPAR

with the highest number of encounters (42%) observed in September. Encounter rates ranged from 0.25 to 0.51 vessels encountered per day per whale during August–October compared to <0.07 vessels per day in all other months in this study. To better inform conservation strategies, further research is required to assess bowhead whale behavioral responses relative to distance from vessels.

KEYWORDS
acoustic disturbance, anthropogenic noise, automatic identification system, bio-logging tags, cetacean, Northwest Passage, vessel strike, vessel traffic

1 | INTRODUCTION

There are two distinct populations of bowhead whales (*Balaena mysticetus*) in the Canadian Arctic: the Bering-Chukchi-Beaufort (BCB) in the west, and the Eastern Canada-West Greenland (ECWG) population in the east (Rugh et al., 2002). The Committee on the Status of Endangered Wildlife in Canada (COSEWIC) has assessed the BCB and ECWG populations to be of Special Concern (Committee on the Status of Endangered Wildlife in Canada, 2009; Fisheries and Oceans Canada, 2014). Both populations were heavily impacted by commercial whaling and are slowly recovering to pre-exploitation numbers (Frasier et al., 2015). Bowhead whales are a culturally important species to Inuit communities in the Canadian Arctic and in West Greenland, where limited and regulated subsistence hunts contribute to traditional practices including food security (Nunavut Wildlife Management Board, 2000; Reeves & Lee, 2022; Suydam & George, 2021).

Bowhead whales are considered highly vulnerable to vessel traffic due to the overlap of their seasonal distribution with Arctic vessel sea routes (Figure 1; Halliday et al., 2021, 2022a,b; Hauser et al., 2018; Martin et al. 2023). In the eastern Canadian Arctic, bowhead whale annual migrations follow the oscillation of sea ice formation and retreat (Fortune et al., 2020a). During winter, ECWG bowhead whales are found in Hudson Strait, northern Hudson Bay, eastern Baffin Island, and the ice edge off West Greenland (Koski et al., 2006; Reeves & Heide-Jørgensen, 1996). In spring, whales from this population reside along the west coast of Greenland, Cumberland Sound, Foxe Basin and Lancaster Sound (Ferguson et al., 2010; Pomerleau et al., 2011). During summer, ECWG bowhead whales are found in bays and fjords in the Canadian High Arctic, such as the Gulf of Boothia, Foxe Basin, eastern Baffin Island, and as far south as Hudson Bay (Cosens & Innes, 2000; Cosens et al., 1997; Ferguson et al., 2010; Higdon & Ferguson, 2010; Pomerleau et al., 2011). Bowhead whales feed on zooplankton (e.g., calanoid copepods; Fortune et al., 2020b,c; Pontbriand et al., 2023), primarily during late summer through autumn (e.g., Finley, 2001; Pomerleau et al., 2011, 2012). However, stable isotope analysis and movement data from tagged whales have revealed that bowhead whales likely feed year-round, although at a lower rate in winter (Fortune et al., 2023; Matthews & Ferguson, 2015; Pomerleau et al., 2018). In autumn, ECWG bowhead whales are found along the east coast of Baffin Island, the west coast of Greenland, and Foxe Basin (Fortune et al., 2023; Reeves & Mitchell, 1990; Reeves et al., 1983). Due to their migration and seasonal movement patterns, ECWG bowhead whales potentially encounter a high number of vessels and associated underwater vessel noise annually, particularly during the open-water season in late summer through autumn (Figure 1; Halliday et al., 2022b).

Economic, geopolitical, and environmental changes resulting from a rapidly warming Arctic climate (Council of Canadian Academies, 2019; Intergovernmental Panel on Climate Change., 2022), has led to a tripling in vessel traffic in the Canadian Arctic over the past three decades (Dawson et al., 2018; Intergovernmental Panel on Climate

17487692, 2024, 4, Downloaded from https://onlinelibrary.wiley.com/doi/10.1111/mms.13125 by Northwest Provincial, Wiley Online Library on [11/10/2024]. See the Terms and Conditions (https://onlinelibrary.wiley.com/terms-and-conditions) on Wiley Online Library for rules of use; OA articles are governed by the applicable Creative Commons License



FIGURE 1 (A) Map of the satellite locations of the 36 telemetry-tagged Eastern Canada-West Greenland (ECWG) bowhead whales during 2012–2017. Whale locations are color coded by year. (B) Map of the vessel track-lines derived from Automatic Identification System (AIS) data during 2012–2017. Vessel track-lines are color coded by year.

Change., 2022). Continued increases in vessel traffic are expected as Arctic shipping routes become ice-free (Bennett et al., 2020; Mudryk et al., 2021; Zeng et al., 2020). Increased vessel traffic can negatively impact bowhead whales in a variety of ways including through an increased risk of collision with vessels, oiling events, and disruption of behavior through vessel presence or noise they generate (George et al., 2017; Halliday, 2020a; Protection of the Arctic Marine Environment, 2019).

Although vessel strike risk and entanglement are presently lower for bowhead whales than for temperate large whale populations residing along heavily developed coastal regions, bowhead whale behavior and increasing vessel traffic due to sea ice reduction elevate these threats (Huntington et al., 2020; Reeves et al., 2012). Bowhead whales are expected to be as vulnerable to vessel strikes as their close relative, the critically endangered North Atlantic right whale *(Eubalaena glacialis)*, because they share similar body morphology and frequently spend time performing surface behaviors, including skim feeding, socializing, nursing, and mating (Dombroski et al., 2021; Laist et al., 2001).

Bowhead whales are continuous ram filter feeders, and may experience reduced feeding efficiency if baleen plates are contaminated by petroleum or become entangled in fishing gear or marine debris (Lambertson et al., 2005). A vertical shift in bowhead prey seasonally occurs during spring and early summer, where aggregations of copepods are found near the surface (Fortune et al., 2023) grazing on phytoplankton. The shallow concentration of prey elevates the risk of vessel strikes. Furthermore, vessel noise is mostly emitted at low frequencies (i.e., <1 kHz) where hearing sensitivity of bowhead whales is inferred to be the most acute (Erbe, 2002; Ketten et al., 2016), although vessels do produce broadband noise that can reach frequencies >10 kHz (Veirs et al., 2016). The presence of vessels and vessel noise has elicited avoidance behavior in some bowhead whales at distances up to 15 km, which can lead to displacement from biologically important areas (Koski & Johnson, 1987; Richardson & Malme, 1993; Richardson et al., 1985a,b; Wartzok et al., 1989). Recent studies have begun to examine the potential overlap or exposure between some Arctic marine mammal populations and vessels or vessel noise

17487692, 2024, 4, Downloaded from https://onlinelibrary.wiley.com/doi/10.1111/mms.13125 by Northwest Provincial, Wiley Online Library on [11/10/2024]. See the Terms and Conditions (https://onlinelibrary.wiley.com/terms-and-conditions) on Wiley Online Library for rules of use; OA articles are governed by the applicable Creative Commons License

(e.g., Halliday, 2020b; Halliday et al., 2018, 2021, 2022a,b; Hauser et al., 2018; Martin et al., 2022, 2023; Reeves et al., 2014). However, a quantitative assessment of the degree of exposure or overlap is needed for effective management.

The current study reports ECWG bowhead whale location data from satellite-linked time and depth transmitters (i.e., bio-logging tags) relative to vessel locations derived from satellite Automatic Identification System (AIS) data. Combining spatial–temporal information about whale and vessel location, we summarize vessel encounter rates for individual whales. The scope of this study encompasses ECWG bowhead whale encounters with vessels from 2012 to 2017 over their range throughout the Eastern Canadian Arctic.

2 | MATERIALS AND METHODS

2.1 | Study area and focal animals

This study focuses on the Eastern Canadian Arctic and West Greenland region, encompassing northern Hudson Bay, Hudson Strait, Foxe Basin, Gulf of Boothia, and Baffin Bay to the west coast of Greenland (Figure 1). During 2012, 2013, and 2016, bowhead whales from the ECWG population were tagged in Cumberland Sound and Foxe Basin, Nunavut (Table S1). Methods for tag deployment are detailed in Ferguson et al. (2010) and Fortune et al. (2020a).

Whales were instrumented with Wildlife Computers location-only SPOT tags, or SPLASH Mk10 tags, which provided information on diving behavior in addition to location data (Wildlife Computers Inc., Redmond, WA; Table S1). Location data were obtained via Doppler effect and the Argos satellite system for the tagged bowhead whales during 2012–2017 (Figure 1a).

Argos satellite tags have been widely used to track large-scale animal movements; however, these tags provide relatively coarse quality locations typically with an accuracy of several hundred meters to several kilometers (https://www.argos-system.org). Each Argos location is assigned a location accuracy designated by a number or letter dependent on the number of orbiting satellites via which the transmission is received, among other factors (https://www.argos-system.org; accuracy categories: 3 [<250 m], 2 [250–500 m], 1 [500–1,500 m], 0 [>1,500 m], A and B [unbounded], Z [invalid]). Consideration of these location accuracies is important when examining fine scale movements of individuals. To increase tag longevity, tags were programmed to limit transmissions to <400 per day during summer, and to <100 transmissions every second day during winter. Depending on satellite reception, programming, and animal behavior (i.e., a tag can only transmit while the animal is at the surface), this transmission rate could lead to temporal gaps of days between satellite-derived locations.

We modeled bowhead locations to account for accuracy issues associated with Argos data using a continuous-time Correlated Random Walk (CRW) model developed by Johnson et al. (2008) and implemented in the "crawl" package (version 2.2/1; Johnson & London, 2018) in R (R Core Team, 2021). The CRW algorithm performs poorly when Argos error greatly exceeds measured values and CRW model priors. Consequently, extreme outliers were removed by filtering the raw location data through the *sdafilter* in the R package "argosfilter" (Freitas et al., 2008; Freitas, 2012). We used the default velocity threshold of 2 m/s, which is identical to the method used by Fortune et al. (2020a) that fit a hierarchical switching-state-space model for the satellite locations from the same tagged individuals. The filter can also be specified to remove outliers which create acute angles in the path of movement (i.e., "spikes"). We used default values for specifying the angular components of the filter; specifically, we removed values that formed angles <15° when they were >2.5 km from the previous location and angles <25° when they were >5 km from the prior location.

The CRW model treats movement as a velocity process with two parameters, β, the autocorrelation in velocity and σ, the variation in velocity. Location error was assumed to be normally distributed with a mean of 0 and a standard deviation equal to that declared by the system operator, Collecte Localisation Satellites, for least-squares location classes 3, 2, and 1 (Collecte Localisation Satellites, 2016). We treated error for the remaining three location

17487692, 2024, 4, Downloaded from https://onlinelibrary.wiley.com/doi/10.1111/mms.13125 by Northwest Provincial, Wiley Online Library on [11/10/2024]. See the Terms and Conditions (https://onlinelibrary.wiley.com/terms-and-conditions) on Wiley Online Library for rules of use; OA articles are governed by the applicable Creative Commons License

classes as parameters to be estimated and fit them to half normal distributions with semi-informative priors. Locations with classes 0, A, and B should have more error than those with a class of 1 ($SD = 1{,}500$ m). Hence, our half normal distributions had a lower bound of 1,500 m. Using data from Vincent et al. (2002), our priors had a mean error of 1,500 m and a standard deviation of 5,000 m for location classes 0 and A, and 7,500 m for location class B. We also set a Laplace prior (double exponential) for β and σ. The Laplace prior had a mean of 3 and a variance of 0.5 on a natural log scale, which is approximately the value of β and σ observed for most species. Note that this is only significant when tracks have few locations. This is the same model and error estimation used for Bering-Chukchi-Beaufort (BCB) bowhead whales in Citta et al. (2018) and Martin et al. (2023). We used the model to better estimate actual bowhead whale locations relative to Argos locations. Whale locations were not regularly spaced in time, and we did not use the model to predict where a whale might be located between Argos locations.

2.2 | Spatial and temporal analysis of ECWG bowhead whales and vessels

Vessel tracks from all months during 2012–2017 were derived from preprocessed satellite AIS data (Spire Global Inc., Cambridge, ON, Canada) in the Eastern Canadian Arctic spanning from Hudson Bay to the west coast of Greenland (Figure 1b), including both class A (mandatory) and class B (voluntary) transmissions. Preprocessing the AIS data involved removing spurious points that were on land, single positions from vessels with no other positions in the vicinity (~100 km), and positions with no maritime mobile service identity (MMSI) number. These former errors are related to either improperly installed AIS equipment or human manipulation to mask true positions of vessels (Jankowski et al., 2021). Positions from AIS data are otherwise assumed to be accurate, given that these locations are based on GPS on board the vessel. For example, one ground truthing study found that AIS positions were accurate to within <100 m of the vessel's actual location (Jankowski et al., 2021). AIS transponders transmit signals that provide the geographic coordinates and other information about individual vessels at regular intervals (e.g., every 2–120 s, depending on vessel behavior and context), and these signals can be received by dedicated satellites and land-based receivers. In earlier years (2012–2015), satellite coverage for AIS was lower in the Arctic, thus leaving some gaps between consecutive locations. However, satellite coverage increased after 2015, leaving smaller gaps between locations and often no gaps for vessels with class A AIS. Vessel traffic in the Eastern Canadian Arctic consists of a variety of vessels including bulk carriers, container ships, cruise ships, government icebreakers and research vessels, tankers, military vessels, ferries, fishing vessels, recreational vessels (i.e., pleasure craft), and local community boats (Arctic Council, 2009; Pizzolato et al., 2016; see Halliday et al., 2022b for a summary).

Several steps were taken to calculate the number of unique vessels that encountered individual tagged whales as well as the closest point of approach (CPA) for each encounter. We used satellite AIS data to calculate the number of unique vessels transiting in the same area (<125 km) and time (<1 hr time differential) as individual tagged whales. We acknowledge this may be a conservative estimate considering that additional vessels may have been present but not transmitting AIS data (i.e., note that only vessels >300 T or with 12 or more passengers are legally required to use AIS, which is class A AIS). Consequently, it is possible that small (~8 m) recreational vessels and local boats are likely underrepresented; however, with the decreasing price of AIS transponders, more recreational vessels are utilizing the technology, particularly in the Arctic where it can be an important safety feature. One vessel class not included in the data set is local boats, which are mostly used for subsistence activities and only range within tens of kilometers from communities and are not typically in the offshore areas preferred by bowhead whales. When a vessel track overlapped in time and space with a tagged bowhead whale, AIS data were used to calculate the range of distances between the vessel and the whale including the CPA during encounters.

A spatial and temporal analysis was completed in ArcGIS using ArcMap 10.8 (ESRI, 2011). First, a buffer radius of 125 km was created around each individual CRW predicted whale location. We selected a 125 km radius because previous modeling suggested that vessel noise can be greater than ambient levels at distances over 100 km in the region (Halliday et al., 2017). Vessel noise can theoretically contribute to ambient sound >125 km, although it is

17487692, 2024, 4, Downloaded from https://onlinelibrary.wiley.com/doi/10.1111/mms.13125 by Northwest Provincial, Wiley Online Library on [11/10/2024]. See the Terms and Conditions (https://onlinelibrary.wiley.com/terms-and-conditions) on Wiley Online Library for rules of use; OA articles are governed by the applicable Creative Commons License

difficult to quantify at this distance (Aulanier et al., 2017). Next, we separated whale location and AIS vessel location data by "day" (i.e., 24-hr period) and paired these data sets in space and time. Individual whale and vessel data from the same day occurring within a radius of 125 km were then extracted and designated as being potentially coincident in the encounter zone. The maximum allowed time differential between a paired whale and vessel location was 1 hr, and the average (± *SD*) time differential was 24.5 (± 17.8) min, with a median of 22.5 min. Vessel speed was not accounted for when pairing locations, therefore, calculated distances between whale and vessel locations represent a snapshot in time. This process was completed separately for each tagged whale to ensure that all possible encounter zones between whales and vessels were included in the analysis. Next, the derived paired whale and vessel locations within each encounter zone were sorted by time to generate a time series of consecutive potential encounters while retaining all underlying data and geographic positional information. The "Points to Line" tool in ArcMap 10.8 was used to calculate the distance (meters) between the closest whale and vessel location as they approached each other within a given encounter zone. For each potential encounter occurring within a radius of 125 km on a given day, the CPA was calculated as the shortest distance observed between the whale and vessel.

If multiple vessels came within 125 km of an individual whale during an encounter, the CPAs from each whale-vessel encounter were compiled and sorted by date and time. This also allowed manual removal of duplicate encounters that resulted from an encounter spanning past midnight where it was included twice based on the original subsetting of data by "day." Once duplicates were removed, all encounters were sorted by the CPA distance and summarized by whale ID (Table S1). Potential hotspots for vessel encounters with distances <50 km were assessed by calculating bivariate normal kernel densities using the CPAs from each whale-vessel encounter using the "ks" package in R (Duong, 2019) with the smoother cross validation (SCV) bandwidth selector. This 50 km radius was chosen because other Arctic whales have been shown to exhibit avoidance responses at this range to vessel noise (Finley et al., 1990; Martin et al., 2022). We defined utilization distributions as the 5%, 25%, 50%, 75%, and 95% probability contours, representing concentration areas (Börger et al., 2006).

2.3 | Ethics approval

This research was conducted in accordance with the Marine Mammal Regulations of the Canadian Fisheries Act. Bowhead whale research was approved under the Fisheries and Oceans Canada License to Fish for Scientific Purposes Nos. S-12/13–1024-NU, S-13/14-1009-NU and S-16/17 1005-NU, and Animal Care Protocols FWI-ACC-3013-018, FWI-ACC-2014-011, FWI-ACC-2016-009.

3 | RESULTS

In the Eastern Canadian Arctic, 36 bowhead whales were tagged during 2012 ($n = 18$), 2013 ($n = 9$), and 2016 ($n = 9$), 19 of which were tagged in Cumberland Sound and 17 in Foxe Basin (see details in Fortune et al., 2020a; Table S1). The sample was composed of 18 females, 16 males, and two whales of unknown sex. Most whales ($n = 28$, 78%) were considered immature based on visually estimated body length (≤13 m), seven whales (19%) were adults (>13 m), and one whale was of an unknown length (Table S1). All bowhead whales in this study were tagged in July or August, which influenced the overall number of encounters (Table S2).

Vessel traffic varied between months and years between 2012 and 2017 (Table 1, Figure 2), such that vessel traffic was always higher between July and November than between December and June, often by nearly an order of magnitude for the number of unique vessels in the study region and by more than an order of magnitude for the distance traveled by all vessels (Table 1). For example, in January to June and December of 2012, 12 unique vessels were present, and they traveled a total distance of 8,482 km, whereas between July and November of the same year, 95 unique vessels were present and they traveled 438,604 km. The number of vessels present and distance traveled

17487692, 2024, 4, Downloaded from https://onlinelibrary.wiley.com/doi/10.1111/mms.13125 by Northwest Provincial, Wiley Online Library on [11/10/2024]. See the Terms and Conditions (https://onlinelibrary.wiley.com/terms-and-conditions) on Wiley Online Library for rules of use; OA articles are governed by the applicable Creative Commons License

TABLE 1 Summary statistics of vessel traffic traveling in the areas of tagged Eastern Canada-West Greenland bowhead whales between 2012 and 2017 when tags were transmitting locations. The area of interest for these statistics is roughly 200 km offshore the east side of Baffin Island up to Devon Island in the north, across to Somerset Island in the west, and south to include all waters of Hudson Bay and Hudson Strait. Months are grouped into the core shipping season (July to November) and the offseason (January to June and December). *N* unique vessels is the number of unique vessels present in the area. See Figure 2 for a display of these same variables by month.

Year	Season	*n* unique vessels	Distance traveled (km)
2012	July–November	95	438,604
	January–June, December	12	8,482
2013	July–November	105	537,229
	January–June, December	15	11,102
2014	July–November	111	556,817
	January–June, December	7	8,945
2015	July–November	113	513,729
	January–June, December	7	9,710
2016	July–November	113	603,778
	January–June, December	18	19,700
2017	July–November	156	685,957
	January–June, December	14	24,812

also generally increased across years, such that in 2017, 156 vessels were present in July to November and they traveled 685,957 km. In every year, the highest number of vessels and distance traveled was in August and September, and the lowest were between January and May (Figure 2), typically with ≤3 vessels per month compared to 57–119 vessels present per month in August and September.

3.1 | Summary of bowhead whale encounters with vessels

During 2012–2017, there were a total of 1,145 encounters within a radius of 125 km among 33 of the 36 tagged whales and tracked vessels (Tables S1 and S2, Figure 3). Three whales (ID numbers: 94542, 114494, and 128149) were never exposed to vessels within this distance radius and were not considered further (Table S1). The 33 exposed whales transmitted location data for variable periods (range: 12–737 days; mean ± *SD*: 303 ± 251 days), with 12 whales transmitting data for over 1 year (Table S1). The number of encounters with AIS vessels for these whales ranged from 1 to 88 per year.

Encounter rate (ER) was scaled to account for the number of days within each month that contained individual whale location transmissions. Whale ID 114500 was tagged in 2012 in Foxe Basin and had the highest encounter rate of all tagged whales (Table S2). Between January and December 2013, this whale encountered 88 AIS vessels, including 65 encounters in September alone, for a daily rate of 2.2 vessels (Table S2). Whale ID 114499 was also tagged in 2012 in Foxe Basin and had the second highest number of encounters with vessels per annum with 87 encounters, during July to December 2012, including 42 encounters in the month of October (ER = 1.4 vessels/day) (Table S2). Not accounting for irregular tagging dates, the month of September in the Eastern Canadian Arctic contained 41.8% of the encounters with vessels ($n = 478$) in this study (Table S2). August and October also contained high numbers of encounters ($n = 199$ and $n = 231$, respectively) between vessels and tagged bowhead whales (Table S2). August, September, and October had the overall highest encounter rates with vessels, with a range of

17487692, 2024, 4, Downloaded from https://onlinelibrary.wiley.com/doi/10.1111/mms.13125 by Northwest Provincial, Wiley Online Library on [11/10/2024]. See the Terms and Conditions (https://onlinelibrary.wiley.com/terms-and-conditions) on Wiley Online Library for rules of use; OA articles are governed by the applicable Creative Commons License



FIGURE 2 Number of unique vessels (top) and the cumulative distance traveled by all vessels (bottom) in each month between January 2012 and December 2017 in the areas used by satellite tagged Eastern Canada-West Greenland bowhead whales. The area of interest for these statistics is roughly 200 km offshore the east side of Baffin Island up to Devon Island in the north, across to Somerset Island in the west, and south to include all waters of Hudson Bay and Hudson Strait.

0.25–0.51 vessels encountered per day compared to <0.07 vessels per day in all other months in this study (Table S2). April ($n = 9$), May ($n = 1$), and June ($n = 14$) had the lowest numbers of encounters and encounter rates annually; however, the overall number of vessel encounters were generally low during November through July compared to August through October for this time series (Table S2). During this time series, a minimum of one whale encountered at least one vessel in each month of the year (Table S2).

The average (± *SD*) CPA between a tagged whale and a vessel was 73.1 (± 31.1) km; however, there were 30 individual whales that had CPAs to vessels at distances ≤50 km for a total of 306 unique encounters (Table S1, Figure 3b). The closest CPA to a vessel was estimated at approximately 2.5 km, and the largest number of

17487692, 2024, 4, Downloaded from https://onlinelibrary.wiley.com/doi/10.1111/mms.13125 by Northwest Provincial, Wiley Online Library on [11/10/2024]. See the Terms and Conditions (https://onlinelibrary.wiley.com/terms-and-conditions) on Wiley Online Library for rules of use; OA articles are governed by the applicable Creative Commons License



FIGURE 3 (A) Map of the 1,145 closest points of approach (CPA) for all encounters between telemetry-tagged Eastern Canada-West Greenland (ECWG) bowhead whales and vessels within a radius of 125 km during 2012–2017. CPAs are color coded by year. (B) Map of the distribution of the 306 CPAs for encounters between tagged ECWG bowhead whales and vessels within a radius of 50 km during 2012–2017. Utilization distributions, calculated via kernel densities, are provided for the 5th, 25th, 50th, 75th, and 95th percentiles.

encounters per year for a single whale at distances ≤50 km was 29 (Table S1). The areas of highest concentrations of ECWG bowhead encounters with vessels at distances ≤50 km were near the two tagging sites, Cumberland Sound and near Igloolik in Foxe Basin, with a third high concentration area in the western Gulf of Boothia (Figure 3b). Additional concentrated distributions of CPA locations at distances ≤50 km were in Hudson Strait, near the community of Qikiqtarjuaq, and in Prince Regent Inlet (Figure 3b).

4 | DISCUSSION

This study summarizes the number of known vessel encounters for tagged ECWG bowhead whales during 2012–2017 in the Eastern Canadian Arctic. Consistently across years, the majority (79.3%) of encounters occurred during August through October (Table S2). Tagged bowhead whales occurred within 3 km from AIS detected vessels. This result differs from the other two species of endemic Arctic cetaceans, beluga whales (*Delphinapterus leucas*) and narwhals (*Monodon monoceros*), which may exhibit strong avoidance reactions to vessels at greater distances (13–50 km; Finley et al., 1990; Martin et al., 2022). However, it is possible certain avoidance behaviors (e.g., vertical avoidance; Senigaglia et al., 2016) occurred and were not captured within the resolution of our data set. This demonstrates the importance of understanding species-specific and context-dependent behavioral responses to vessels for effective management and conservation (see Gomez et al., 2016 for a review). Further research is needed to examine individual bowhead whale behavioral responses, including dive responses, age/sex-related responses, and the potential for habituation or sensitization to vessels at a range of distances. If bowhead whales fail to exhibit an avoidance reaction to vessels, this could place them at higher risk of vessel strikes (Halliday et al., 2022b).

17487692, 2024, 4, Downloaded from https://onlinelibrary.wiley.com/doi/10.1111/mms.13125 by Northwest Provincial, Wiley Online Library on [11/10/2024]. See the Terms and Conditions (https://onlinelibrary.wiley.com/terms-and-conditions) on Wiley Online Library for rules of use; OA articles are governed by the applicable Creative Commons License

Richardson and Malme (1993) described bowhead whale avoidance behavior as an increase in swimming speed away from the vessel with a decrease in dive duration and number of blows per surfacing. Avoidance responses in bowhead whales have been documented at distances <1–4 km and rarely beyond 8 km from standard vessels when underway (Richardson & Malme, 1993; Richardson et al., 1985a,b). Bowhead whale avoidance responses to seismic survey vessels have been reported at distances between 20 and 30 km (Koski & Johnson, 1987; Miller et al., 1997, 1999; Richardson et al., 1999). Studies suggest bowhead whales that are socializing or foraging appear more tolerant of potential sources of anthropogenic disturbance than migrating whales (Koski et al., 2009; Miller et al., 2005; Quakenbush et al., 2010; Robertson et al., 2013; Wartzok et al., 1989). This information is important in the interpretation of the results from this study, which indicate that the highest concentrations of ECWG bowhead encounters with vessels at ≤50 km distance occurred when whales are presumed to be feeding during August–October near Igloolik in Foxe Basin, the Gulf of Boothia, and Cumberland Sound (Figure 3b). During our time series in the Gulf of Boothia, government vessels were the most frequent vessel class, followed by tanker ships (Halliday et al., 2022b). Vessel traffic in Cumberland Sound is comprised of low levels of bulk carriers, cruise ships, government vessels, and recreational boats, and very low levels of ferries and tanker ships (Halliday et al., 2022b). These areas host large annual congregations of bowhead whales which promote more social encounters and serve as feeding grounds and migratory corridors during the late summer through autumn seasons (Fortune et al., 2020a; Pomerleau et al., 2011).

A modeling study by Halliday et al. (2022b) reported the highest relative vessel strike risk areas for ECWG bowhead whales were the Gulf of Boothia, Cumberland Sound, and near Isabella Bay (eastern coast of Baffin Island), Nunavut, Canada. Vessel strike risk was highest in August and September, corresponding with monthly levels of vessel traffic (Halliday et al., 2022b). Our study and Halliday et al. (2022b) used some of the same input data; however, Halliday et al. (2022b) included data prior to 2012 and from whales tagged in West Greenland. The inclusion of whales from West Greenland showed that Isabella Bay could be a potential hotspot for overlap between bowhead whales and vessel traffic, because whales from West Greenland tend to spend their summers foraging near Isabella Bay (Chambault et al., 2018). Conversely, whales from this study, which were tagged in Foxe Basin and Cumberland Sound, did not use Isabella Bay. We therefore stress the need for caution when interpreting our results, because our study did not include whales tagged in West Greenland as those tag data (2001–2011) do not overlap in time with the satellite AIS vessel tracking data (post 2011). Isabella Bay appears to be an important area for a segment of the population of bowhead whales during the summer, which was not captured by the data used in our study and warrants further study. Regardless, the collective findings of our study and Halliday et al. (2022b) highlight the need for vessel management within these high use areas (Figure 3b) particularly during late summer and autumn to reduce the potential for vessel strikes and noise disturbance.

There are other emerging threats to bowhead whales that are also important to consider in addition to impacts from increasing vessel traffic in the Eastern Canadian Arctic. Climate change has led to increased predation risk from killer whales (*Orcinus orca*) (Breed, 2021; Matthews et al., 2020), and the potential for thermal stress from increased sea temperatures (Chambault et al., 2018). Furthermore, changes have been identified in zooplankton species composition and abundance as an index of ocean productivity and prey quality with more lipid-poor copepod species recently found in bowhead whale habitats (Fortune et al., 2020b; Møller & Nielsen, 2020). These cumulative threats may impact the species' predation risk, habitat use patterns, and foraging success which may ultimately affect the health of individuals and survival of the species at the population level.

There are caveats associated with this study that need to be considered. These include (1) the inherent spatial accuracy in some whale location data derived from the Argos satellite system; however, the CRW model accounts to some degree for issues associated with spatial accuracy. Many encounters between bowhead whales and vessels at distances ≤50 km could not be investigated further primarily due to insufficient tag location data. This resulted in the inability to model the encounters to identify any specific bowhead whale behavioral responses to vessels. (2) The whales tagged in this study may not represent the full population due to sample size and potential differences in behavior based on individual differences, specific areas, and seasons (i.e., spatial segregation based on age and sex;

17487692, 2024, 4, Downloaded from https://onlinelibrary.wiley.com/doi/10.1111/mms.13125 by Northwest Provincial, Wiley Online Library on [11/10/2024]. See the Terms and Conditions (https://onlinelibrary.wiley.com/terms-and-conditions) on Wiley Online Library for rules of use; OA articles are governed by the applicable Creative Commons License

Heide-Jørgensen et al., 2010). For example, the lack of data from Isabella Bay, as mentioned previously. (3) Two of the regions with the highest concentrations of whale encounters with vessels at distances ≤50 km predominantly align with the two tagging sites, Cumberland Sound and near Igloolik in Foxe Basin (Figure 3). It is plausible that these concentrations were influenced by the proximity of telemetry locations to the tagging sites, indicating a potential spatial bias in the recorded encounters. (4) It is possible that there were vessels present which did not carry AIS transponders. Due to these limitations, our results underestimate bowhead whale encounters with vessels within the 125-km radius in the Eastern Canadian Arctic.

Given the cultural and ecological importance of this vulnerable species, this summary of encounters provides evidence that further studies are needed, possibly including controlled experiments, to better understand how bowhead whales may respond to vessels or vessel noise at known distances and across behavioral states. Future studies also should include focused recordings of underwater noise unique to each vessel and vessel class, including measurements of source levels and frequency spectra. With developments in tag technology, future incorporation of 3-dimensional movement and acoustic recordings would provide the opportunity to examine changes in the acoustic behavior of bowhead whales, identify the acoustic signature, received level, and exact time when vessel noise is received at the whale, and ultimately allow an in-depth examination of bowhead whale behavioral response to the type and received level of vessel noise and other sounds. Similarly, long-term tag technology has also advanced, and the use of tags equipped with Fastloc GPS and higher resolution dive profiles (e.g., Martin et al., 2022) could yield more data on reactions of bowhead whales to vessels.

ACKNOWLEDGMENTS

We are grateful for the invaluable logistical support provided by the Hunters and Trappers Organizations in Igloolik, Sanirajak, and Pangnirtung. Field assistance and vessel operations were provided by Levi Qaunaq and Natalino Piugattak from Igloolik, and Noah Ishulutaq and Timeosie Akpalialuk from Pangnirtung, and Bernard LeBlanc for leading the tagging and biopsy collection. Satellite AIS data were provided by the Marine Environmental Observation Prediction and Response (MEOPAR) network through partnerships with Spire Global Inc. (formerly exactEarth Ltd.). We thank Dr. John Citta for sharing R code for the CRW model.

AUTHOR CONTRIBUTIONS

Morgan Jennifer Martin: Conceptualization; formal analysis; funding acquisition; investigation; methodology; project administration; software; visualization; writing – original draft; writing – review and editing. **William David Halliday:** Conceptualization; data curation; funding acquisition; methodology; project administration; software; supervision; validation; writing – review and editing. **Steven Ferguson:** Conceptualization; data curation; funding acquisition; resources; writing – review and editing. **Brent G Young:** Data curation; writing – review and editing. **Rachel Charish:** Formal analysis; software; writing – review and editing. **Jackie Dawson:** Data curation; writing – review and editing. **Sarah M.E. Fortune:** Data curation; methodology; writing – review and editing. **Francis Juanes:** Supervision; writing – review and editing. **Levi Qaunaq:** Data curation; resources. **Steve J. Insley:** Conceptualization; funding acquisition; project administration; supervision; writing – review and editing.

COMPETING INTERESTS

The authors declare there are no competing interests.

DATA AVAILABILITY

Data generated or analyzed during this study are available from the corresponding author upon reasonable request.

ORCID

Morgan J. Martin https://orcid.org/0000-0002-3556-6632
William D. Halliday https://orcid.org/0000-0001-7135-076X

17487692, 2024, 4, Downloaded from https://onlinelibrary.wiley.com/doi/10.1111/mms.13125 by Northwest Provincial, Wiley Online Library on [11/10/2024]. See the Terms and Conditions (https://onlinelibrary.wiley.com/terms-and-conditions) on Wiley Online Library for rules of use; OA articles are governed by the applicable Creative Commons License

Steven H. Ferguson https://orcid.org/0000-0002-3794-0122
Francis Juanes https://orcid.org/0000-0001-7397-0014

REFERENCES

Arctic Council. (2009, April). *Arctic marine shipping assessment report*.

Aulanier, F., Simard, Y., Roy, N., Gervaise, C., & Bandet, M. (2017). Effects of shipping on marine acoustic habitats in Canadian Arctic estimated via probabilistic modeling and mapping. *Marine Pollution Bulletin*, *125*(1–2), 115–131.

Bennett, M. M., Stephenson, S. R., Yang, K., Bravo, M. T., & De Jonghe, B. (2020). The opening of the Transpolar Sea Route: Logistical, geopolitical, environmental, and socioeconomic impacts. *Marine Policy*, *121*, Article 104178.

Börger, L., Franconi, N., De Michele, G., Gantz, A., Meschi, F., Manica, A., Lovari, S., & Coulson, T. (2006). Effects of sampling regime on the mean and variance of home range size estimates. *Journal of Animal Ecology*, *75*, 1393–1405. https://doi.org/10.1111/j.1365-2656.2006.01164.x

Breed, G. A. (2021). Predators and impacts of predation. In J. C. George & J. G. M. Thewissen (Eds.), The bowhead whale *Balaena mysticetus:* Biology and human interactions (pp. 457–70). Academic Press.

Chambault, P., Albertsen, C. A., Patterson, T. A., Hansen, R. G., Tervo, O., Laidre, K. L., & Heide-Jørgensen, M. P. (2018). Sea surface temperature predicts the movements of an Arctic cetacean: the bowhead whale. *Scientific Reports*, *8*, Article 9658. https://doi.org/10.1038/s41598-018-27966-1

Citta, J. J., Lowry, L. F., Quakenbush, L. T., Kelly, B. P., Fischbach, A. S., London, J. M., Jay, C. V., Frost, K. J., O'Corry-Crowe, G., Crawford, J. A., Boveng, P. L., Cameron, M., Von Duyke, A. L., Nelson, M., Harwood, L. A., Richard, P., Suydam, R., Heide-Jørgensen, M. P., Hobbs, R. C., … Gray, T. (2018). A multi-species synthesis of satellite telemetry data in the Pacific Arctic (1987–2015): Overlap of marine mammal distributions and core use areas. *Deep Sea Research Part II: Topical Studies in Oceanography*, *152*, 132–153. https://doi.org/10.1016/j.dsr2.2018.02.006

Collecte Localisation Satellites. (2016). Argos user's manual. https://d9-wret.s3.us-west-2.amazonaws.com/assets/palladium/production/s3fs-public/atoms/files/Argos_users_manual_en_2016_v1.6.6-r363_9.pdf

Committee on the Status of Endangered Wildlife in Canada. (2009). *COSEWIC assessment and update status report on the Bowhead Whale Balaena mysticetus, Bering–Chukchi–Beaufort population and Eastern Canada–West Greenland population, in Canada*.

Cosens, S. E., & Innes, S. (2000). Distribution and numbers of bowhead whales (*Balaena mysticetus*) in northwestern Hudson Bay in August 1995. *Arctic*, *53*(1), 36–41. https://doi.org/10.14430/arctic832

Cosens, S. E., Qamukaq, T., Parker, B., Dueck, L. P., & Anardjuak, B. (1997). The distribution and numbers of Bowhead whales, *Balaena mysticetus*, in northern Foxe Basin in 1994. *Canadian Field-Naturalist*, *111*, 381–388.

Council of Canadian Academies. (2019). *Canada's top climate change risks*. The Expert Panel on Climate Change Risks and Adaptation Potential, Ottawa, Canada.

Dawson, J., Pizzolato, L., Howell, S. E., Copland, L., & Johnston, M. E. (2018). Temporal and spatial patterns of ship traffic in the Canadian Arctic from 1990 to 2015. *Arctic*, *71*(1), 15–26. https://doi.org/10.14430/arctic4698

Dombroski, J. R., Parks, S. E., & Nowacek, D. P. (2021). Dive behavior of North Atlantic right whales on the calving ground in the Southeast USA: implications for conservation. *Endangered Species Research*, *46*, 35–48. https://doi.org/10.3354/esr01141

Duong, T. (2019). *Ks: Kernel smoothing* (R package version 1.11.5) [Computer software]. https://cran.r-hub.io/web/packages/ks/ks.pdf

Erbe, C. (2002). Hearing abilities of baleen whales. *Defence Research and Development Canada Atlantic CR* 2002–065.

ESRI. (2011). ArcGIS desktop: release 10 [Computer software]. Environmental Systems Research Institute, Redlands, CA.

Ferguson, S. H., Dueck, L., Loseto, L. L., & Luque, S. P. (2010). Bowhead whale *Balaena mysticetus* seasonal selection of sea ice. *Marine Ecology Progress Series*, *411*, 285–297. https://doi.org/10.3354/meps08652

Finley, K. J. (2001). Natural history and conservation of the Greenland whale, or bowhead, in the Northwest Atlantic. *Arctic*, *54*(1), 55–76. https://doi.org/10.14430/arctic764

Finley, K. J., Miller, G. W., Davis, R. A., & Greene, C. R., Jr. (1990). Reactions of belugas (*Delphinapterus leucas*) and narwhals (*Monodon monoceros*) to ice-breaking ships in the Canadian High Arctic. *Canadian Bulletin of Fisheries and Aquatic Sciences*, *224*, 97–117.

Fisheries and Oceans Canada. (2014). Management Plan for the Bering-Chukchi-Beaufort population of bowhead whale (Balaena mysticetus) in Canada. Species at Risk Act Management Plan Series. Fisheries and Oceans Canada. https://www.canada.ca/en/environment-climate-change/services/species-risk-public-registry/management-plans/bowhead-whale-bering-chukchi-beaufort.html

Fortune, S. M., Young, B. G., & Ferguson, S. H. (2020a). Age-and sex-specific movement, behaviour and habitat-use patterns of bowhead whales (*Balaena mysticetus*) in the Eastern Canadian Arctic. *Polar Biology*, *43*, 1725–1744.

Fortune, S. M., Ferguson, S. H., Trites, A. W., Hudson, J. M., & Baumgartner, M. F. (2020b). Bowhead whales use two foraging strategies in response to fine-scale differences in zooplankton vertical distribution. *Scientific Reports*, *10*, Article 20249. https://doi.org/10.1038/s41598-020-76071-9

17487692, 2024, 4, Downloaded from https://onlinelibrary.wiley.com/doi/10.1111/mms.13125 by Northwest Provincial, Wiley Online Library on [11/10/2024]. See the Terms and Conditions (https://onlinelibrary.wiley.com/terms-and-conditions) on Wiley Online Library for rules of use; OA articles are governed by the applicable Creative Commons License

Fortune, S. M. E., Ferguson, S. H., Trites, A. W., Leblanc, B., Lemay, V., Hudson, J. M., & Baumgartner, M. F. (2020c). Seasonal diving and foraging behaviour of eastern Canada-West Greenland bowhead whales. *Marine Ecology Progress Series*, *643*, 197–217. https://doi.org/10.3354/meps13356

Fortune, S. M. E., Trites, A. W., LeMay, V., Baumgartner, M. F., & Ferguson, S. H. (2023). Year-round feeding on lipid-rich copepods across large spatial scales suggest that bowhead whales have the potential to adapt to climate change. *Frontiers in Marine Science*, *9*. https://doi.org/10.3389/fmars.2022.853525

Frasier, T. R., Petersen, S. D., Postma, L., Johnson, L., Heide-Jørgensen, M. P., & Ferguson, S. H. (2015). Abundance estimates of the Eastern Canada-West Greenland bowhead whale *(Balaena mysticetus)* population based on genetic capture-mark-recapture analyses. Canadian Science Advisory Secretariat Research Document.

Freitas, C. (2012). *argosfilter: Argos locations filter* (R package version 0.63) [Computer software]. R Foundation for Statistical Computing. https://CRAN.R-project.org/package=argosfilter

Freitas, C., Lydersen, C., Fedak, M. A., & Kovacs, K. M. (2008). A simple new algorithm to filter marine mammal Argos locations. *Marine Mammal Science*, *24*(2), 315–325. https://doi.org/10.1111/j.1748-7692.2007.00180.x

George, J. C., Sheffield, G., Reed, D. J., Tudor, B., Stimmelmayr, R., Person, B. T., Sformo, T., & Suydam, R. (2017). Frequency of injuries from line entanglements, killer whales, and ship strikes on Bering-Chukchi-Beaufort Seas bowhead whales. *Arctic*, *70*(1), 37–46. https://doi.org/10.14430/arctic4631

Gomez, C., Lawson, J. W., Wright, A. J., Buren, A. D., Tollit, D., & Lesage, V. (2016). A systematic review on the behavioural responses of wild marine mammals to noise: the disparity between science and policy. *Canadian Journal of Zoology*, *94*(12), 801–819. https://doi.org/10.1139/cjz-2016-0098

Halliday, W. D. (2020a). Literature review of ship strike risk to whales (Report written for Fisheries and Oceans Canada). Wildlife Conservation Society, Toronto, Canada. https://library.wcs.org/Scientific-Research/Research-Publications/Publications-Library/ctl/view/mid/40093/pubid/DMX3964300000.aspx

Halliday, W. D. (2020b). Vessel traffic trends in the arctic and overlap with important marine mammal areas *(Report for Transport Canada)*. Wildlife Conservation Society, Toronto, Canada. https://library.wcs.org/Scientific-Research/Research-Publications/Publications-Library/ctl/view/mid/40093/pubid/DMX3964000000.aspx

Halliday, W. D., Dawson, J., Yurkowski, D. J., Doniol-Valcroze, T., Ferguson, S. H., Gjerdrum, C., Hussey, N. E., Kochanowicz, Z., Mallory, M. L., Marcoux, M., & Watt, C. A. (2022a). Vessel risks to marine wildlife in the Tallurutiup Imanga National Marine Conservation Area and the eastern entrance to the Northwest Passage. *Environmental Science & Policy*, *127*(80), 181–195. https://doi.org/10.1016/j.envsci.2021.10.026

Halliday, W. D., Insley, S. J., Hilliard, R. C., de Jong, T., & Pine, M. K. (2017). Potential impacts of shipping noise on marine mammals in the western Canadian Arctic. *Marine Pollution Bulletin*, *123*, 73–82. https://doi.org/10.1016/j.marpolbul.2017.09.027

Halliday, W. D., Le Baron, N., Citta, J., Dawson, J., Doniol-Valcroze, T., Ferguson, M., Ferguson, S. H., Fortune, S., Harwood, L. A., Heide-Jørgensen, M. P., Lea, E., Quakenbush, L., Young, B., Yurkowski, D., & Insley, S. J. (2022b). Overlap between bowhead whales (Balaena mysticetus) and vessel traffic in the North American Arctic and implications for conservation and management. *Biological Conservation*, *276*, Article 109820. https://doi.org/10.1016/j.biocon.2022.109820

Halliday, W. D., Pine, M. K., Citta, J. J., Harwood, L., Hauser, D. D. W., Hilliard, R. C., Lea, E. V., Loseto, L. L., Quakenbush, L., & Insley, S. J. (2021). Potential exposure of beluga and bowhead whales to underwater noise from ship traffic in the Beaufort and Chukchi Seas. *Ocean & Coastal Management*, *204*, Article 105473. https://doi.org/10.1016/j.ocecoaman.2020.105473

Halliday, W. D., Têtu, P.-L., Dawson, J., Insley, S. J., & Hilliard, R. C. (2018). Tourist vessel traffic in important whale areas in the western Canadian Arctic: risks and possible management solutions. *Marine Policy*, *97*, 72–81. https://doi.org/10.1016/j.marpol.2018.08.035

Hauser, D. D. W., Laidre, K. L., & Stern, H. L. (2018). Vulnerability of Arctic marine mammals to vessel traffic in the increasingly ice-free Northwest Passage and Northern Sea Route. *Proceedings of the National Academy of Sciences of the United States of America*, *115*, 7617–7622. https://doi.org/10.1073/pnas.1803543115

Heide-Jørgensen, M., Laidre, K., Wiig, Ø., Postma, L., Dueck, L., & Bachmann, L. (2010). Large-scale sexual segregation of bowhead whales. *Endangered Species Research*, *13*(1), 73–78. https://doi.org/10.3354/esr00315

Higdon, J. W., & Ferguson, S. H. (2010). Past, present, and future for bowhead whales (*Balaena mysticetus*) in northwest Hudson Bay. In S. H. Ferguson, L. L. Loseto, & M. L. Mallory (Eds.), *A little less Arctic: top predators in the world's largest northern inland sea, Hudson Bay* (pp. 159–177). Springer.

Huntington, H. P., Danielson, S. L., Wiese, F. K., Baker, M., Boveng, P., Citta, J. J., De Robertis, A., Dickson, D. M. S., Farley, E., George, J. C., Iken, K., Kimmel, D. G., Kuletz, K., Ladd, C., Levine, R., Quakenbush, L., Stabeno, P., Stafford, K. M., Stockwell, D., & Wilson, C. (2020). Evidence suggests potential transformation of the Pacific Arctic Ecosystem is underway. *Nature Climate Change*, *10*, 342–348. https://doi.org/10.1038/s41558-020-0695-2

17487692, 2024, 4, Downloaded from https://onlinelibrary.wiley.com/doi/10.1111/mms.13125 by Northwest Provincial, Wiley Online Library on [11/10/2024]. See the Terms and Conditions (https://onlinelibrary.wiley.com/terms-and-conditions) on Wiley Online Library for rules of use; OA articles are governed by the applicable Creative Commons License

Intergovernmental Panel on Climate Change. (2022). Climate change 2022: Impacts, adaptation and vulnerability. In H.-O. Pörtner, D. C. Roberts, M. Tignor, E. S. Poloczanska, K. Mintenbeck, A. Alegría, M. Craig, S. Langsdorf, S. Löschke, V. Möller, A. Okem, & B. Rama (Eds.), *Contribution of Working Group II to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change*. Cambridge University Press. https://doi:10.1017/9781009325844

Jankowski, D., Lamm, A., & Hahn, A. (2021). Determination of AIS position accuracy and evaluation of reconstruction methods for maritime observation data. *IFAC-PapersOnLine*, *54*(16), 97–104. https://doi.org/10.1016/j.ifacol.2021.10.079

Johnson, D. S., & London, J. M. (2018). *R package 'crawl'*. [Computer software]. https://cran.r-project.org/web/packages/crawl/

Johnson, D. S., London, J. M., Lea, M. A., & Durban, J. W. (2008). Continuous-time correlated random walk model for animal telemetry data. *Ecology*, *89*(5), 1208–1215. https://doi.org/10.1890/07-1032.1

Ketten, D. R., Arruda, J., Cramer, S., & Yamato, M. (2016). Great ears: Low-frequency sensitivity correlates in land and marine leviathans. In A. N. Popper & A. Hawkins (Eds.), *The effects of noise on aquatic life II* (pp. 529–538). Springer.

Koski, W. R., Funk, D. W., Ireland, D. S., Lyons, C., Christie, K., Macrander, A. M., & Blackwell, S. B. (2009). *An update on feeding by bowhead whales near an offshore seismic survey in the central Beaufort Sea* (SC/61/BRG3). International Whaling Commission. http://iwc.int/index.php?cID=1780&cType=document

Koski, W. R., Heide-Jørgensen, M. P., & Laidre, K. L. (2006). Winter abundance of bowhead whales, *Balaena mysticetus*, in the Hudson Strait, March 1981. *Journal of Cetacean Research and Management*, *8*(2), 139–144. https://doi.org/10.47536/jcrm.v8i2.710

Koski, W. R., & Johnson, S. R. (1987). *Behavioral studies and aerial photogrammetry*. In Responses of bowhead whales to an offshore drilling operation in the Alaskan Beaufort Sea, autumn 1986 (Sect. 4). Report from LGL Limited, King City, Ontario, and Greeneridge Sciences Inc., Santa Barbara, CA, for Shell Western Exploration and Production Inc., Anchorage, AK.

Laist, D. W., Knowlton, A. R., Mead, J. G., Collet, A. S., & Podesta, M. (2001). Collisions between ships and whales. *Marine Mammal Science*, *17*(1), 35–75. https://doi.org/10.1111/j.1748-7692.2001.tb00980.x

Lambertson, R. H., Rasmussen, K. J., Lancaster, W. C., & Hintz, R. J. (2005). Functional morphology of the mouth of the bowhead whale and its implications for conservation. *Journal of Mammalogy*, *86*, 342–352. https://doi.org/10.1644/ber-123.1

Martin, M. J., Halliday, W. D., Citta, J., Quakenbush, L., Harwood, L., Lea, E. V., Juanes, F., Dawson, J., Nicoll, A., & Insley, S. J. (2023). Exposure and behavioral responses of tagged bowhead whales (*Balaena mysticetus*) to vessels in the Pacific Arctic. *Arctic Science*, *9*(3), 600–615. https://doi.org/10.1139/as-2022-0052

Martin, M. J., Halliday, W. D., Storrie, L., Citta, J., Dawson, J., Hussey, N. E., Juanes, F., Loseto, L. L., MacPhee, S. A., Moore, L., Nicoll, A., O'Corry-Crowe, G., & Insley, S. J. (2022). Exposure and behavioral responses of tagged beluga whales (*Delphinapterus leucas*) to ships in the Pacific Arctic. *Marine Mammal Science*, *39*(2), 387–421. https://doi.org/10.1111/mms.12978

Matthews, C. J., Breed, G. A., LeBlanc, B., & Ferguson, S. H. (2020). Killer whale presence drives bowhead whale selection for sea ice in Arctic seascapes of fear. *Proceedings of the National Academy of Sciences of the United States of America*, *117*(12), 6590–6598. https://doi.org/10.1073/pnas.1911761117

Matthews, C. J. D., & Ferguson, S. H. (2015). Seasonal foraging behaviour of eastern Canada-West Greenland bowhead whales: An assessment of isotopic cycles along baleen. *Marine Ecology Progress Series*, *522*, 269–286. https://doi.org/10.3354/meps11145

Miller, G. W., Elliot, R. E., Koski, W. R., & Richardson, W. J. (1997). *Northstar marine mammal monitoring program, 1996: Marine mammal and acoustical monitoring of a seismic program in the Alaskan Beaufort Sea* (pp. 5.1–5.115.). Report from LGL Limited, King City, ON, to BP Exploration (Alaska) Inc., Anchorage, AK.

Miller, G. W., Elliot, R. E., Koski, W. R., & Richardson, W. J. (1999). Whales. *Marine Mammal and Acoustical Monitoring Program of Western Geophysical's Open-Water Seismic Program in the Alaskan Beaufort Sea, 1998: 90-Day Report* (pp. 5.1–5.59). Report from LGL Limited, King City, ON, to Western Geophysical, Houston TX and National Marine Fisheries Service, Anchorage, AK and Silver Spring, MD.

Miller, G. W., Moulton, V. D., Davis, R. A., Holst, M., Millman, P., MacGillivray, A., & Hannay, D. (2005). Monitoring seismic effects on marine mammals-southeastern Beaufort Sea, 2001–2002. In S. L. Armsworthy, P. J. Cranford, & K. Lee (Eds.), *Offshore oil and gas environmental effects monitoring/approaches and technologies* (pp. 511–542). Battelle Press.

Møller, E. F., & Nielsen, T. G. (2020). Borealization of Arctic zooplankton—smaller and less fat zooplankton species in Disko Bay, Western Greenland. *Limnology and Oceanography*, *65*(6), 1175–1188. https://doi.org/10.1002/lno.11380

Mudryk, L. R., Dawson, J., Howell, S. E., Derksen, C., Zagon, T. A., & Brady, M. (2021). Impact of 1, 2 and 4 °C of global warming on ship navigation in the Canadian Arctic. *Nature Climate Change*, *11*, 673–679. https://doi.org/10.1038/s41558-021-01087-6

Nunavut Wildlife Management Board. (2000). *Final report of the Inuit bowhead whale study, Nunavut*, Canada. Iqaluit, NU.

17487692, 2024, 4, Downloaded from https://onlinelibrary.wiley.com/doi/10.1111/mms.13125 by Northwest Provincial, Wiley Online Library on [11/10/2024]. See the Terms and Conditions (https://onlinelibrary.wiley.com/terms-and-conditions) on Wiley Online Library for rules of use; OA articles are governed by the applicable Creative Commons License

Pizzolato, L., Howell, S. E., Dawson, J., Laliberté, F., & Copland, L. (2016). The influence of declining sea ice on shipping activity in the Canadian Arctic. *Geophysical Research Letters*, *43*(23), 12–146. https://doi.org/10.1002/2016GL071489

Pomerleau, C., Lesage, V., Ferguson, S. H., Winkler, G., Petersen, S. D., & Higdon, J. W. (2012). Prey assemblage isotopic variability as a tool for assessing diet and the spatial distribution of bowhead whale *Balaena mysticetus* foraging in the Canadian eastern Arctic. *Marine Ecology Progress Series*, *469*, 161–174.

Pomerleau, C., Matthews, C. J. D., Gobeil, C., Stern, G. A., Ferguson, S. H., & Macdonald, R. W. (2018). Mercury and stable isotope cycles in baleen plates are consistent with year-round feeding in two bowhead whale (*Balaena mysticetus*) populations. *Polar Biology*, *41*(5), 1881–1893.

Pomerleau, C., Patterson, T., Luque, S., Lesage, V., Heide-Jørgensen, M., Dueck, L., & Ferguson, S. (2011). Bowhead whale *Balaena mysticetus* diving and movement patterns in the eastern Canadian Arctic: implications for foraging ecology. *Endangered Species Research*, *15*(2), 167–177. https://doi.org/10.3354/esr00373

Pontbriand, T., Davoren, G. K., Fortune, S. M. E., Pomerleau, C., Young, B. G., & Ferguson, S. H. (2023). Differences in individual bowhead whale *Balaena mysticetus* habitat use, foraging dive depth and diet during the peak feeding season. *Marine Ecology Progress Series*, *706*, 109–123. https://doi.org/10.3354/meps14238

Protection of the Arctic Marine Environment. (2019). *Underwater noise in the Arctic: A state of knowledge report*. PAME Secretariat, Akureyri, Iceland.

Quakenbush, L. T., Small, R. J., & Citta, J. J. (2010). *Satellite tracking of western Arctic bowhead whales* (OCS Study BOEM 2013-01110). U.S. Department of the Interior, Bureau of Ocean Energy Management, Alaska Outer Continental Shelf Region, Anchorage, AK.

R Core Team. (2021). R: A language and environment for statistical computing *(Version 4.1.1)* [Computer software]. R Foundation for Statistical Computing.

Reeves, R., Mitchell, E., Mansfield, A., & McLaughlin, M. (1983). Distribution and migration of the bowhead whale, *Balaena mysticetus*, in the Eastern North American Arctic. *Arctic*, *36*, 5–64.

Reeves, R., Rosa, C., George, J. C., Sheffield, G., & Moore, M. (2012). Implications of Arctic industrial growth and strategies to mitigate future vessel and fishing gear impacts on bowhead whales. *Marine Policy*, *36*, 454–462. https://doi.org/10.1016/j.marpol.2011.08.005

Reeves, R. R., Ewins, P. J., Agbayani, S., Heide-Jørgensen, M. P., Kovacs, K. M., Lydersen, C., Suydam, R., Elliott, W., Polet, G., van Dijk, Y., & Blijleven, R. (2014). Distribution of endemic cetaceans in relation to hydrocarbon development and commercial shipping in a warming Arctic. *Marine Policy*, *44*, 375–389. https://doi.org/10.1016/j.marpol.2013.10.005

Reeves, R. R., & Heide-Jørgensen, M. P. (1996). Recent status of bowhead whales, *Balaena mysticetus*, in the wintering grounds off West Greenland. *Polar Research*, *15*(2), 115–125. https://doi.org/10.3402/POLAR.V15I2.6641

Reeves, R., & Lee, D. S. (2022). Bowhead whales and whaling in the central and eastern Canadian Arctic, 1970–2021. *Journal of Cetacean Research and Management*, *23*(1), 1–25. https://doi.org/10.47536/jcrm.v23i1.359

Reeves, R. R., & Mitchell, E. (1990). Bowhead whales in Hudson Bay, Hudson Strait, and Foxe Basin: A review. *Naturaliste canadien*, *117*, 25–43.

Richardson, W. J., Fraker, M. A., Würsig, B., & Wells, R. S. (1985a). Behaviour of bowhead whales *Balaena mysticetus* summering in the Beaufort Sea: Reactions to industrial activities. *Biological Conservation*, *32*(3), 195–230. https://doi.org/10.1016/0006-3207(85)90111-9

Richardson, W. J., & Malme, C. I. (1993). *Man-made noise and behavioral responses*. In J. J. Burns, J. J. Montague, & C. J. Cowles (Eds.), *The bowhead whale* (pp. 631–700). Special Publication Number 2, Society for Marine Mammalogy.

Richardson, W. J., Miller, G. W., & Greene, C. R., Jr. (1999). Displacement of migrating bowhead whales by sounds from seismic surveys in shallow waters of the Beaufort Sea. *Journal of the Acoustical Society of America*, *106*(4_Supplement), Article 2281. https://doi.org/10.1121/1.427801

Richardson, W. J., Wells, R. S., & Würsig, B. (1985b). Disturbance responses of bowheads, 1980–84. In W. J. Richardson (Ed.), Behavior, disturbance responses and distribution of bowhead whales *Balaena mysticetus* in the eastern Beaufort Sea, 1980–84 (pp. 89–196). Report from LGL Ecological Research Associates Inc., Bryan, TX, for U.S. Minerals Management Service. NTIS No. PB87-124376.

Robertson, F. C., Koski, W. R., Thomas, T. A., Richardson, W. J., Würsig, B., & Trites, A. W. (2013). Seismic operations have variable effects on dive-cycle behavior of bowhead whales in the Beaufort Sea. *Endangered Species Research*, *21*(2), 143–160. https://doi.org/10.3354/esr00515

Rugh, D. J., Demaster, D., Rooney, A., Breiwick, J. M., Shelden, K., & Moore, S. E. (2002). A review of bowhead whale (*Balaena mysticetus*) stock identity. *Journal of Cetacean Research and Management*, *5*(3), 267–279. https://doi.org/10.47536/jcrm.v5i3.807

Senigaglia, V., Christiansen, F., Bejder, L., Gendron, D., Lundquist, D., Noren, D.P., Schaffar, A., Smith, J. C., Williams, R., Martinez, E., Stockin, K., & Lusseau, D. (2016). Meta-analyses of whale-watching impact studies: Comparisons of cetacean responses to disturbance. *Marine Ecology Progress Series*, *542*, 251–263. https://doi.org/10.3354/MEPS11497

17487692, 2024, 4, Downloaded from https://onlinelibrary.wiley.com/doi/10.1111/mms.13125 by Northwest Provincial, Wiley Online Library on [11/10/2024]. See the Terms and Conditions (https://onlinelibrary.wiley.com/terms-and-conditions) on Wiley Online Library for rules of use; OA articles are governed by the applicable Creative Commons License

Suydam, R., & George, J. C. (2021). Current indigenous whaling. In J. C. George & J. G. M. Thewissen (Eds.), *The bowhead whale* (pp. 519–535). Elsevier. https://doi.org/10.1016/B978-0-12-818969-6.00032-7

Veirs, S., Veirs, V., & Wood, J. D. (2016). Ship noise extends to frequencies used for echolocation by endangered killer whales. *PeerJ*, *4*(3), e1657. https://doi.org/10.7717/peerj.1657

Vincent, C., McConnell, B. J., Ridoux, V., & Fedak, M. A. (2002). Assessment of Argos location accuracy from satellite tags deployed on captive gray seals. *Marine Mammal Science*, *18*(1), 156–166. https://doi.org/10.1111/j.1748-7692.2002.tb01025.x

Wartzok, D., Watkins, W. A., Würsig, B., & Malme, C. I. (1989). *Movements and behaviors of bowhead whales in response to repeated exposures to noises associated with industrial activities in the Beaufort Sea*. Report from Purdue University, Fort Wayne, IN, for Amoco Production Co., Denver, CO.

Zeng, Q., Lu, T., Lin, K.-C., Yuen, K. F., & Li, K. X. (2020). The competitiveness of Arctic shipping over Suez Canal and China-Europe railway. *Transport Policy*, *86*, 34–43. https://doi.org/10.1016/j.tranpol.2019.11.005

SUPPORTING INFORMATION

Additional supporting information can be found online in the Supporting Information section at the end of this article.

How to cite this article: Martin, M. J., Halliday, W. D., Ferguson, S. H., Young, B. G., Charish, R., Dawson, J., Fortune, S. M. E., Juanes, F., Qaunaq, L., & Insley, S. J. (2024). Exposure of satellite tagged bowhead whales (*Balaena mysticetus*) to transiting vessels in the Eastern Canadian Arctic. *Marine Mammal Science*, *40*(4), e13125. https://doi.org/10.1111/mms.13125

17487692, 2024, 4, Downloaded from https://onlinelibrary.wiley.com/doi/10.1111/mms.13125 by Northwest Provincial, Wiley Online Library on [11/10/2024]. See the Terms and Conditions (https://onlinelibrary.wiley.com/terms-and-conditions) on Wiley Online Library for rules of use; OA articles are governed by the applicable Creative Commons License